

UNIQUENESS OF UNCONDITIONAL BASES IN c_0 -PRODUCTS

P.G. CASAZZA AND N.J. KALTON

ABSTRACT. We give counterexamples to a conjecture of Bourgain, Casazza, Lindenstrauss and Tzafriri that if X has a unique unconditional basis (up to permutation) then $c_0(X)$ also has a unique unconditional basis. We also give some positive results including a simpler proof that $c_0(\ell_1)$ has a unique unconditional basis and a proof that $c_0(\ell_{p_n}^{N_n})$ has a unique unconditional basis when $p_n \downarrow 1$, $N_{n+1} \geq 2N_n$ and $(p_n - p_{n+1}) \log N_n$ remains bounded.

1. INTRODUCTION

A Banach space X is said to have a *unique unconditional basis* (or more precisely, a unique unconditional basis up to permutation) if it has an unconditional basis and if whenever (u_n) and (v_n) are two normalized unconditional bases of X , then there is permutation π of $\mathbb{N}$ such that (v_n) and $(u_{\pi(n)})$ are equivalent. Since unconditional bases correspond to discrete or atomic order-continuous lattice structures on X , this can be reworded as a statement that such a lattice-structure is essentially unique.

The earliest examples of Banach spaces with unique unconditional bases are c_0, ℓ_1 ([10]) and ℓ_2 ([9]). It was shown by Lindenstrauss and Zippin [12] that amongst spaces with symmetric bases this is the complete list. Later Edelstein and Wojtaszczyk showed that direct sums of these spaces also have unique unconditional bases. All these results can be found in [11]. In [3] the authors attempted a complete classification and showed that the spaces $c_0(\ell_1), c_0(\ell_2), \ell_1(c_0)$ and $\ell_1(\ell_2)$ all have unique unconditional bases while $\ell_2(\ell_1)$ does not. They also found an unexpected additional space, 2-convexified Tsirelson (see [5] for the definition) with a unique unconditional basis. Recently, the authors found a new

1991 *Mathematics Subject Classification.* 46B15, 46B07.

The first author was supported by NSF Grant 9706108; the second author was supported by Grant DMS-9500125

approach to this type of problem and were able to add some more spaces, including Tsirelson space (see [5]) itself and certain Nakano spaces [4] (as pointed out in [4] some spaces considered by Gowers [8] provide further examples) ; we also showed that uniqueness of the unconditional basis need not be inherited by a complemented subspace.

This note is motivated by a question raised in [3]. They asked if $c_0(X)$ has a unique unconditional basis whenever X does. The idea here is that if this and the corresponding dual result for ℓ_1 -products holds then one could iterate the results in [3] to produce examples such as $c_0(\ell_1(c_0(\ell_1)))$ and so on.

Unfortunately as we show below in Section 4, the answer to this question is negative and Tsirelson space T or its 2-convexified version both produce counterexamples. However, we show how our approach in [4] can be used for c_0 -products. We give a much shorter proof (Theorem 3.3) of the fact that $c_0(\ell_1)$ has a unique unconditional basis; the original proof of this result in [3] is extremely technical. We show by the same techniques (Theorem 3.4) that examples of the type $c_0(\ell_{p_n}^{N_n})$ where $p_n \downarrow 1$, $N_{n+1} \geq 2N_n$ and $(p_n - p_{n+1}) \log N_n$ remains bounded must also have unique unconditional bases.

In Section 4, we also use the same techniques to show that for certain right-dominant spaces X , as introduced in [4], such as Tsirelson space T , any unconditional basis of $c_0(X)$ must be equivalent to a *subset* of the canonical basis (Theorem 4.1). Nevertheless we show that the unconditional basis of $c_0(T)$ is *not* unique as already remarked.

We conclude this section with a few remarks on terminology and assumptions. We will frequently index unconditional bases and basic sequences by an unordered countable index set $\mathcal{N}$ which need not be the natural numbers $\mathbb{N}$. We will assume that any unconditional basic sequence $(u_n)_{n \in \mathcal{N}}$ is semi-normalized, i.e. $0 < \inf_{n \in \mathcal{N}} \|u_n\| \leq \sup_{n \in \mathcal{N}} \|u_n\| < \infty$. We will say that two unconditional basic sequences $(u_n)_{n \in \mathcal{N}}$ and $(v_n)_{n \in \mathcal{N}'}$ are *equivalent* if there is a bijection $\pi : \mathcal{N} \rightarrow \mathcal{N}'$ so that $(u_n)_{n \in \mathcal{N}}$ and $(v_{\pi(n)})_{n \in \mathcal{N}}$ are equivalent.

An unconditional basic sequence $(u_n)_{n \in \mathcal{N}}$ in X is *complemented* if there is a bounded projection $P : X \rightarrow [u_n]_{n \in \mathcal{N}}$. If $(u_n)_{n \in \mathcal{N}}$ is an unconditional basis of X and $(v_n)_{n \in \mathcal{N}'}$ is an unconditional basic sequence of the form $v_n = \sum_{k \in \mathcal{A}_n} a_k u_k$ where the sets $(\mathcal{A}_n)_{n \in \mathcal{N}'}$ are disjoint subsets of $\mathcal{N}$ we say that $(v_n)_{n \in \mathcal{N}'}$ is *disjoint* with respect to $(u_n)_{n \in \mathcal{N}}$. If (v_n) is a complemented disjoint sequence then it may be shown that there is a projection $Px = \sum_{n \in \mathcal{N}'} v_n^*(x) v_n$ where each $v_n^* \in X^*$

is of the form $v_n^* = \sum_{k \in \mathcal{A}_n} b_k u_k^*$ where $(u_k^*)_{k \in \mathcal{N}}$ are the sequence of biorthogonal functions for $(u_k)_{k \in \mathcal{N}}$.

It will be convenient to represent a space X with unconditional basis $(u_n)_{n \in \mathcal{N}}$ as a sequence space modelled on the index set $\mathcal{N}$, identifying $\sum_{k \in \mathcal{N}} a_k u_k$ with the function $f : \mathcal{N} \rightarrow \mathbb{R}$ given by $f(k) = a_k$. This identifies X as a discrete Banach lattice and allows us to use functional notation. The canonical basis of a sequence space X modelled on $\mathcal{N}$ is denoted by $(e_n)_{n \in \mathcal{N}}$.

If $(u_n)_{n \in \mathcal{N}}$ is an unconditional basis for X and N is a natural number we denote by $(u_n)_{n \in \mathcal{N}}^N$ the naturally induced unconditional basis of X^N (the direct sum of N copies of X).

For future reference we note here that our techniques depend critically on the following result, proved in Theorem 3.5 of [4]:

Theorem 1.1. *Suppose X is a Banach space with an unconditional basis $(u_n)_{n \in \mathcal{N}}$ which does not contain uniformly complemented copies of ℓ_2^n (i.e. is not sufficiently Euclidean). Suppose $(v_n)_{n \in \mathcal{B}}$ is a complemented unconditional basic sequence in X . Then there is a integer N and a complemented disjoint sequence $(w_n)_{n \in \mathcal{B}}$ in the basis $(u_n)_{n \in \mathcal{N}}^N$ such that $(v_n)_{n \in \mathcal{B}}$ is equivalent to $(w_n)_{n \in \mathcal{B}}$.*

2. A CRITERION FOR AN ℓ_1 - OR c_0 -PRODUCT TO BE SUFFICIENTLY EUCLIDEAN

The aim of this section is to establish criteria for c_0 -product to contain uniformly complemented copies of ℓ_2^n so that we can apply Theorem 1.1.

If X is a Banach space we will say that X has property $P(k, M)$ where $k \in \mathbb{N}$ and $M \geq 1$ if whenever $S : \ell_2^k \rightarrow X$ and $T : X \rightarrow \ell_2^k$ are operators satisfying $TS = I_{\ell_2^k}$ then $\|S\| \|T\| \geq M$. We will say that a sequence of Banach spaces $(X_j)_{j=1}^\infty$ has property $P(k, M)$ if each X_j has property $P(k, M)$. A Banach space X (respectively a sequence of Banach spaces $(X_j)_{j=1}^\infty$) is *sufficiently Euclidean* if there exists M so that X (respectively $(X_j)_{j=1}^\infty$) fails $P(k, M)$ for every $k \in \mathbb{N}$.

We recall that if H is a finite-dimensional Hilbert space and $A : H \rightarrow X$ is any linear map then the ℓ -norm of A is given by:

$$\ell(A) = \mathbf{E}(\|\sum_{i=1}^m g_i A e_i\|^2)^{1/2}$$

where $(e, \dots, e_m)$ is any orthonormal basis of H and $(g_1, \dots, g_m)$ is a sequence of independent normalized Gaussian random variables. See [15]. If S is an operator

on a Banach space X and E is a closed subspace of X we denote by S_E the restriction of S to E .

Lemma 2.1. *There exists a universal constant $c > 0$ with the following property. Suppose H is an n -dimensional Hilbert space and X is any Banach space. Suppose $S : H \rightarrow X$ is any operator with $\|S\| \leq 1$. Then there is a subspace E of H with $\dim E \geq c\ell(S)^2$ so that $\|S_E\| \leq 3\ell(S)n^{-1/2}$.*

Proof. It will suffice to prove this for S one-to-one, since the result then follows by a simple perturbation argument. Let μ be normalized invariant measure on the surface of the sphere in ℓ_2^n . Consider the norm $\xi \rightarrow \|S\xi\|$; this satisfies $\|S\xi\| \leq \|\xi\|$ for all ξ . We use Theorem 4.2 of [13] (p.12). If M_r is a median value of the norm $\|S\xi\|$ then

$$M_r \leq \sqrt{2} \left(\int \|S\xi\|^2 d\mu \right)^{1/2} = \sqrt{2/n} \ell(S).$$

Lemma 2.2. *Suppose X is a Banach space with property $P(k, M)$. Suppose H is an n -dimensional Hilbert space and $S : H \rightarrow X$ and $T : X \rightarrow H$ are bounded operators with $\|T\| \leq 1$. Then*

$$|\operatorname{tr}(TS)| \leq Cn^{1/2} \max(M^{-1}\ell(S), k^{1/2}\|S\|)$$

for some universal constant C .

Proof. Suppose $1 \leq j \leq n$ and that s_j is the j th. singular value of TS . We can restrict to a subspace H_j of dimension j so that $\|TS\xi\| \geq s_j\|\xi\|$ for all $\xi \in H_j$.

Assume $s_j > 0$. Then by the preceding Lemma 2.1 there is a subspace E of H_j so that $\dim E \geq c\ell(S_{H_j})^2$ and $\|S_E\| \leq 3\ell(S_E)j^{-1/2}$.

Suppose $\dim E \geq k$. Then we must have $\|T\|\|S_E\| \geq Ms_j$ so that $\ell(S_E) \geq \frac{1}{3}Ms_jj^{1/2}$. This implies that $s_j \leq 3M\ell(S)j^{-1/2}$. If $\dim E < k$ then

$$\ell(S_{H_j})^2 \|S_{H_j}\|^{-2} \leq c^{-1}k$$

and so $\ell(TS_{H_j}) \leq c^{-1/2}k^{1/2}\|S\|$. From this we deduce that $j^{1/2}s_j \leq c^{-1/2}k^{1/2}\|S\|$ or $s_j \leq c^{-1/2}k^{1/2}j^{-1/2}\|S\|$. Combining we obtain that

$$s_j \leq \max(3M^{-1}\ell(S), c^{-1/2}k^{1/2}\|S\|)j^{-1/2}.$$

Now

$$|\operatorname{tr}(TS)| \leq \sum_{j=1}^n s_j \leq Cn^{1/2} \max(M^{-1}\ell(S), k^{1/2}\|S\|)$$

for some universal constant C .

Lemma 2.3. *There is a universal constant C so that if X is a Banach space with property $P(k, M)$ then whenever H is a Hilbert space of dimension n , and $S : H \rightarrow X$ and $T : X \rightarrow H$ are bounded operators with $\|T\| \leq 1$, we have*

$$|\operatorname{tr}(TS)| \leq C\ell(S)\left(\frac{1}{M} + k^{1/2}(\log n)^{-1/2}\right)n^{1/2}.$$

Proof. We first choose an orthonormal basis $(e_i)_{i=1}^n$ of H so that $\|Se_i\| = \|S_{H_i}\|$ where $H_i = [e_i, e_{i+1}, \dots, e_n]$. Pick $m = \lceil n^{1/2} \rceil$. Then if $g_1, \dots, g_m$ are normalized independent Gaussians,

$$\ell(S) \geq \mathbf{E}\left(\left\|\sum_{i=1}^m g_i T e_i\right\|^2\right)^{1/2} \geq \|Se_m\| \mathbf{E}\left(\max_{1 \leq i \leq m} |g_i|\right).$$

Now (cf. [13] p. 23) this implies that

$$\|Se_m\| \leq C(\log m)^{-1/2}\ell(S)$$

for some universal constant C . Our choice of m implies that we can replace this estimate by

$$\|Se_m\| \leq C(\log n)^{-1/2}\ell(S)$$

for some universal constant C .

If $E = [e_1, \dots, e_m]$ then

$$|\operatorname{tr}(TS_E)| \leq m^{1/2}\ell(S_E) \leq n^{1/4}\ell(S).$$

On the other hand

$$|\operatorname{tr}(TS_{H_{m+1}})| \leq C \max(M^{-1}, k^{1/2}(\log n)^{-1/2})\ell(S)n^{1/2}$$

by Lemma 2.2. Combining these results gives us our estimate. $\square$

Proposition 2.4. *Suppose $(X_j)_{j=1}^\infty$ is not sufficiently Euclidean. Then $\ell_1(X_j)$ is not sufficiently Euclidean.*

Proof. Suppose (X_j) satisfies property $P(k, M)$. Suppose $n \in \mathbb{N}$ and $S : \ell_2^n \rightarrow \ell_1(X_j)$ and $T : \ell_1(X_j) \rightarrow \ell_2^n$ are any operators satisfying $TS = I_{\ell_2^n}$; we assume that $\|T\| = 1$. We write $S\xi = (S_i\xi)_{i=1}^\infty$ and $T(x_i)_{i=1}^\infty = \sum_{i=1}^\infty T_i x_i$.

Now $n = \text{tr} (TS) = \sum_{i=1}^{\infty} \text{tr} (T_i S_i)$. On the other hand, by Lemma 2.3 we have that:

$$(2.1) \quad |\text{tr} (T_i S_i)| \leq C n^{1/2} \left(\frac{1}{M} + k^{1/2} (\log n)^{-1/2} \ell(S_i) \right).$$

Let $(e_1, \dots, e_n)$ be any orthonormal basis. Then by the Kahane-Khintchine inequality we have that

$$(2.2) \quad \ell(S_i) \leq C_0 \mathbf{E} \left(\left\| \sum_{j=1}^n g_j S_i e_j \right\| \right)$$

where the $(g_i)_{i=1}^n$ are normalized independent Gaussians, and C_0 is a universal constant. Hence

$$(2.2) \quad \sum_{i=1}^{\infty} \ell(S_i) \leq C_0 \mathbf{E} \left(\left\| \sum_{j=1}^n g_j S e_j \right\| \right) \leq C \ell(S).$$

Hence, combining (2.1), (2.2) and (2.3), and since $n = \sum_{i=1}^{\infty} \text{tr} (T_i S_i)$,

$$n \leq C_0 n^{1/2} \left(\frac{1}{M} + k^{1/2} (\log n)^{-1/2} \right) \ell(S) \leq C_1 n \|S\| \left(\frac{1}{M} + k^{1/2} (\log n)^{-1/2} \right)$$

for some universal constant C_1 . We thus obtain an estimate

$$\|S\| \geq C_1^{-1} \min(M, k^{-1/2} (\log n)^{1/2})$$

for some absolute constant C . This establishes the result. $\square$

Remark. In the case when $X = c_0$ we have that X satisfies $P(ck^{1/2}, k)$ for $c > 0$ and all k . We thus obtain that $\ell_1(X)$ satisfies $P(c(\log k)^{1/4}, k)$ for some $c > 0$ and all k . On the other hand Figiel, Lindenstrauss and Milman [7] established the upper estimate that $\ell_1(c_0)$ contains a subspace 2-isomorphic to ℓ_2^k which is $(\log k)^{1/2}$ -complemented; this estimate is best possible (see [2]). This suggests our method, while not optimal, cannot be improved significantly.

Corollary 2.5. *Suppose $(X_j)_{j=1}^{\infty}$ is not sufficiently Euclidean. Then $c_0(X_j)$ is not sufficiently Euclidean.*

Proof. This follows by simple duality. $\square$

3. UNCONDITIONAL BASES IN c_0 -PRODUCTS

For convenience we define a sequence space X as a Köthe space of real-valued functions on a countable set I (with counting measure) so that the canonical basis vectors $(e_i)_{i \in I}$ form a 1-unconditional basis. Usually of course we take $I = \mathbb{N}$, but for our purposes it is convenient also to allow $I = \mathbb{N} \times \mathbb{N}$ and certain other alternatives. A typical element x of X is of the form $x = (x(i))_{i \in I}$.

Let $(u_n)_{n \in \mathcal{N}}$ be a set of disjointly supported vectors in X . Then $(u_n)_{n \in \mathcal{N}}$ is an unconditional basic sequence, which is complemented if and only if there exists a biorthogonal sequence $(u_n^*)_{n \in \mathcal{N}} \in X^*$ with $\text{supp } u_n^* \subset \text{supp } u_n$, $u_n u_n^* \geq 0$, $\langle u_n, u_n^* \rangle = 1$ and such that the projection

$$Px = \sum_{n \in \mathcal{N}} \langle x, u_n^* \rangle u_n$$

is well-defined and bounded. If we define $f_n = u_n u_n^*$ then $f_n \geq 0$, $f_n \in \ell_1(I)$ and $\|f_n\|_1 = 1$ for all $n \in \mathbb{N}$. Under these circumstances we say that (u_n) is a complemented disjoint sequence and we assume that (u_n^*) and (f_n) are associated with (u_n) . Note that we can always replace u_n and u_n^* by $|u_n|$ and $|u_n^*|$ and hence also assume them positive.

We start with an observation which we will use repeatedly.

Lemma 3.1. *Suppose X is a sequence space (modelled on an index set I) and $(u_n)_{n \in \mathcal{N}}$ is a complemented disjoint sequence. Let $(A_n)_{n \in \mathcal{N}}$ be any sequence of disjoint sets such that, for some $\delta > 0$ we have $\|f_n \chi_{A_n}\|_1 \geq \delta > 0$ for all $n \in \mathcal{N}$. Let $v_n = u_n \chi_{A_n}$. Then $(v_n)_{n \in \mathcal{N}}$ is a complemented disjoint sequence equivalent to $(u_n)_{n \in \mathcal{N}}$. Furthermore the biorthogonal vectors v_n^* may be chosen so that $v_n v_n^* \leq \delta^{-1} f_n$.*

Proof. Let P be the projection onto $[u_n]_{n \in \mathcal{N}}$ as defined above. Let

$$v_n^* = \|f_n \chi_{A_n}\|_1^{-1} u_n^* \chi_{A_n}$$

and define

$$Qx = \sum_{n \in \mathcal{N}} \langle x, v_n^* \rangle v_n.$$

Then it is easy to verify that Q is a bounded projection, $Qu_n = v_n$ and $P(v_n) = \|f_n \chi_{A_n}\|_1 u_n$. This quickly establishes the equivalence of $(u_n)_{n \in \mathcal{N}}$ and $(v_n)_{n \in \mathcal{N}}$. Furthermore $v_n^* v_n \leq \delta^{-1} f_n$. $\square$

Next suppose $(X_i)_{i=1}^\infty$ is a sequence of sequence spaces modelled on index sets J_i (either finite sets or $\mathbb{N}$). We suppose that for some $q < \infty$ the spaces (X_i) satisfy a lower q -estimate uniformly, i.e. there exists $c > 0$ so that if $i \in \mathbb{N}$ and $x_1 \dots, x_n$ are disjoint in X_i then

$$\left\| \sum_{k=1}^n x_k \right\|_{X_i} \geq c \left(\sum_{k=1}^n \|x_k\|_{X_i}^q \right)^{1/q}.$$

Let $Y = c_0(X_j)$ be the sequence space on $J = \{(i, j) : j \in J_i, i \in \mathbb{N}\}$ of all $x = (x[i])_{i=1}^\infty$, where $x[i] \in X_i$ are so that $\lim_{i \rightarrow \infty} \|x[i]\|_{X_i} = 0$. We define

$$\|x\|_Y = \max_{i \in \mathbb{N}} \|x[i]\|_{X_i}.$$

Now suppose $(u_n)_{n \in \mathcal{N}}$ is a complemented disjoint sequence in Y , with biorthogonal sequence (u_n^*) . As above let $f_n = u_n^* u_n$. Then define $F_n \in \ell_1(\mathbb{N})$ by

$$F_n(i) = \sum_{j \in J_i} f_n(i, j).$$

We will say that (u_n) is C -tempered if

$$\sum_{i=1}^\infty \sup_{n \in \mathcal{N}} F_n(i) \leq C.$$

Theorem 3.2. *Suppose $(X_i)_{i=1}^\infty$ is a sequence of sequence spaces satisfying a uniform lower q -estimate for some $q < \infty$. Suppose $(u_n)_{n \in \mathcal{N}}$ is a normalized complemented disjoint sequence in $c_0(X_i)$. Then there is a complemented disjoint sequence $(v_n)_{n \in \mathcal{N}}$ equivalent to $(u_n)_{n \in \mathcal{N}}$ and a partition $\mathcal{N} = \cup_{n \in \mathcal{A}} \mathcal{B}_n$ of $\mathcal{N}$ with the following properties:*

- (1) *For each $i \in \mathbb{N}$ we have either $\|v_k[i]\|_{X_i} = 0$ or 1.*
- (2) *For some C each $(v_k)_{k \in \mathcal{B}_n}$ is C -tempered.*
- (3) *There exists an integer N and subsets $(S_n)_{n \in \mathcal{A}}$ of $\mathbb{N}$ such that $\sum_{n \in \mathcal{A}} \chi_{S_n} \leq N - 1$ and $v_k[i] = 0$ whenever $k \in \mathcal{B}_n$ and $i \notin S_n$. Hence for any finitely nonzero sequence $(a_n)_{n \in \mathcal{N}}$,*

$$\max_{n \in \mathcal{A}} \left\| \sum_{k \in \mathcal{B}_n} a_k v_k \right\|_Y \leq \left\| \sum_{k \in \mathcal{N}} a_k v_k \right\|_Y \leq (N - 1) \max_{n \in \mathcal{A}} \left\| \sum_{k \in \mathcal{B}_n} a_k v_k \right\|_Y.$$

Proof. As usual we let P be the induced projection on $[u_n]_{n \in \mathcal{N}}$.

First let $A_n = \{(i, j) : \|u_n[i]\|_{X_i} \geq (2\|P\|)^{-1}\}$. Notice that

$$\sum_{i \notin A_n} \langle u_n[i], u_n[i]^* \rangle \leq \frac{1}{2\|P\|} \sum_{i=1}^{\infty} \|u_n[i]\|_{X_i^*} \leq \frac{1}{2\|P\|} \|u_n\|_{Y^*} \leq \frac{1}{2}.$$

Hence $(u_n \chi_{A_n})_{n \in \mathcal{N}}$ is a complemented disjoint sequence equivalent to $(u_n)_{n \in \mathcal{N}}$. It follows after some appropriate renormalization that we can replace $(u_n)_{n \in \mathcal{N}}$ by an equivalent sequence with the additional property that $\|u_n[i]\|_{X_i} = 1$ or $u_n[i] = 0$ for every $i \in \mathbb{N}$. For each $n \in \mathcal{N}$ let $S_n = \{i : \|u_n[i]\|_{X_i} = 1\}$.

Next fix any $N \in \mathbb{N}$ so that $N > 1 + c^{-q}(1 + \|P\|)^q$, where c is the constant of the uniform lower q -estimate. Let $\delta = N^{-1}$. We pick a maximal subset $\mathcal{A}$ of $\mathcal{N}$ with the property that if $\mathcal{F}$ is a subset of $\mathcal{A}$ with $|\mathcal{F}| \leq N$ then

$$\sum_{i=1}^{\infty} \max_{n \in \mathcal{F}} F_n(i) \geq (1 - \delta)|\mathcal{F}|.$$

(Here $F_n(i) = \sum_{j \in J_i} f_n(i, j) = \langle u_n[i], u_n^*[i] \rangle$ as usual.)

Now let $\mathcal{F}$ be any subset of $\mathcal{A}$ with $|\mathcal{F}| = N$. We can partition $\mathbb{N}$ into N disjoint sets $(A_n)_{n=1}^N$ so that if $i \in A_n$ then $F_n(i) = \max_{m \in \mathcal{F}} F_m(i)$. Let $v = \sum_{n \in \mathcal{F}} u_n \chi_{A'_n}$ where $A'_n = \{(i, j) : i \in A_n, j \in J_i\}$. Clearly $\|v\|_Y \leq 1$. However

$$Pv = \sum_{n \in \mathcal{F}} \sum_{i \in A_n} F_n(i) u_n.$$

Since

$$\sum_{n \in \mathcal{F}} \sum_{i \in A_n} F_n(i) \geq N(1 - \delta) = N - 1$$

we conclude that

$$\left\| \sum_{n \in \mathcal{F}} u_n \right\|_Y \leq \|P\| + 1.$$

On the other hand,

$$\left\| \sum_{n \in \mathcal{F}} u_n \right\|_Y \geq c \max_{i \in \mathbb{N}} \left(\sum_{n \in \mathcal{F}} \chi_{S_n}(i) \right)^{1/q}.$$

Hence

$$(3.1) \quad \max_{i \in \mathbb{N}} \sum_{n \in \mathcal{F}} \chi_{S_n}(i) \leq c^{-q}(\|P\| + 1)^q < N.$$

Now suppose for some i we have $\sum_{n \in \mathcal{A}} \chi_{S_n}(i) \geq N$. Then we can find a subset $\mathcal{F}$ of $\mathcal{A}$ with $|\mathcal{F}| = N$ so that $\chi_{S_n}(i) = 1$ for $n \in \mathcal{F}$ contradicting (3.1). We therefore conclude that:

$$\max_{i \in \mathbb{N}} \sum_{n \in \mathcal{A}} \chi_{S_n}(i) \leq N - 1.$$

Now suppose $k \in \mathcal{N} \setminus \mathcal{A}$. There exists a subset $\mathcal{F}$ of $\mathcal{A}$ with $1 \leq |\mathcal{F}| \leq N - 1$ and such that

$$\sum_{i=1}^{\infty} \max(F_k(i), \max_{n \in \mathcal{F}} F_n(i)) < (|\mathcal{F}| + 1)(1 - \delta).$$

Hence

$$\begin{aligned} \sum_{i=1}^{\infty} \min(F_k(i), \max_{n \in \mathcal{F}} F_n(i)) &= \sum_{i=1}^{\infty} (F_k(i) + \max_{n \in \mathcal{F}} F_n(i)) - \sum_{i=1}^{\infty} \max(F_k(i), \max_{n \in \mathcal{F}} F_n(i)) \\ &> \sum_{i=1}^{\infty} F_k(i) - 1 + \delta \\ &> \delta. \end{aligned}$$

Thus there exists $n \in \mathcal{F}$ so that

$$\sum_{i=1}^{\infty} \min(F_k(i), F_n(i)) > \delta/N = \delta^2.$$

Now let $T_k = \{i : F_k(i) < 2N^2 F_n(i)\}$. Then $\sum_{i \in T_k} F_k(i) > \frac{1}{2}\delta^2$. In the case when $k \in \mathcal{A}$ we will put $T_k = S_k$.

We now can partition $\mathcal{N}$ into disjoint sets $(\mathcal{B}_n)_{n \in \mathcal{A}}$ so that $n \in \mathcal{B}_n$ and if $k \in \mathcal{B}_n$ then $\sum_{i \in T_k} F_k(i) > \frac{1}{2}\delta^2$ and $F_k(i) < 2N^2 F_n(i)$ for $i \in T_k \subset S_n$. If we then let $T'_k = \{(i, j) : i \in T_k, j \in J_i\}$ and $v_k = u_k \chi_{T'_k}$ then by Lemma 3.1 we have that $(v_k)_{k \in \mathcal{N}}$ is a complemented disjoint sequence in Y equivalent to $(u_k)_{k \in \mathcal{N}}$. Furthermore if (v_k^*) is the biorthogonal sequence then we have an estimate $v_k v_k^* \leq M f_k$ for a suitable constant M . It follows that for each $n \in \mathcal{A}$ we have

$$\langle v_k[i], v_k^*[i] \rangle \leq 2MN^2 F_n(i)$$

whenever $k \in \mathcal{B}_n$. Thus the sets $(v_k)_{k \in \mathcal{B}_n}$ are each C -tempered where C is a constant depending only on c, q and $\|P\|$.

Finally suppose $(a_n)_{n \in \mathcal{N}}$ is finitely non-zero. Then

$$\begin{aligned}
\left\| \sum_{k \in \mathcal{N}} a_k v_k \right\|_Y &= \max_{i \in \mathbb{N}} \left\| \sum_{k \in \mathcal{N}} a_k v_k[i] \right\|_{X_i} \\
&= \max_{i \in \mathbb{N}} \left\| \sum_{n \in \mathcal{A}, i \in S_n} \sum_{k \in \mathcal{B}_n} a_k v_k[i] \right\|_{X_i} \\
&\leq (N-1) \max_{n \in \mathcal{A}} \left\| \sum_{k \in \mathcal{B}_n} a_k v_k \right\|_Y.
\end{aligned}$$

This completes the proof. $\square$

Let us first use this result to give a simpler proof of the result of [3] that $c_0(\ell_1)$ has a unique unconditional basis (up to permutation).

Theorem 3.3. *The space $c_0(\ell_1)$ has a unique unconditional basis.*

Proof. We start with the remark that $c_0(\ell_1)$ is not sufficiently Euclidean (cf. Bourgain [2] or Corollary 2.5 above). Hence any complemented unconditional basic sequence is equivalent to a complemented positive disjoint sequence in $c_0(\ell_1)^m$ for some m . We can clearly suppose $m = 1$.

We next show that any C -tempered C -complemented disjoint sequence $(u_n)_{n \in \mathcal{N}}$ is K -equivalent to the standard ℓ_1 -basis where K depends only on C . Indeed we may suppose $\|u_n[i]\|_1 = 1$ or $u_n[i] = 0$ for each i, n . Let $G(i) = \max_n \langle u_n[i], u_n^*[i] \rangle$. Then $\|G\|_1 \leq C$.

Now

$$\begin{aligned}
\left\| \sum_{n \in \mathcal{N}} a_n u_n \right\|_{c_0(\ell_1)} &\geq \frac{1}{C} \sum_{i=1}^{\infty} G(i) \sum_{u_n[i] \neq 0} |a_n| \\
&= \frac{1}{C} \sum_{n \in \mathcal{N}} |a_n| \sum_{u_n[i] \neq 0} G(i) \\
&\geq \frac{1}{C} \sum_{n \in \mathcal{N}} |a_n| \sum_{i=1}^{\infty} \langle u_n[i], u_n^*[i] \rangle \\
&= \frac{1}{C} \sum_{n \in \mathcal{N}} |a_n|.
\end{aligned}$$

It follows that any unconditional basis of $c_0(\ell_1)$ is equivalent to the canonical unconditional basis of $c_0(X_n)$ where each X_n is either ℓ_1 or ℓ_1^m for some $m = m(n)$. However there must be infinitely many indices n for which $X_n = \ell_1$ (since $c_0(\ell_1)$ cannot be decomposed as $\ell_1 \oplus Z$ where Z contains no copy of ℓ_1 .) It then easily follows that $c_0(\ell_1)$ has a unique unconditional basis. $\square$

Theorem 3.4. *Suppose $1 < p_n < \infty$ and $p_n \downarrow 1$. Let (N_n) be an increasing sequence of natural numbers such that $(p_n - p_{n+1}) \log N_n$ is bounded and $N_{n+1} \geq 2N_n$. Then $c_0(\ell_{p_n}^{N_n})$ has a unique unconditional basis.*

Proof. We suppose the sequence (p_n) fixed and first consider $c_0(\ell_{p_n}^{M_n})$ for any sequence of integers (M_n) . It is easy to see by considering the ultraproduct $\prod_{\mathcal{U}} \ell_{p_n}^{M_n}$ (which is an L_1 -space) that the sequence $(\ell_{p_n}^{M_n})$ is not sufficiently Euclidean. By Corollary 2.5 the space $Y = c_0(\ell_{p_n}^{M_n})$ is also not sufficiently Euclidean and therefore every complemented unconditional basic sequence in Y is equivalent to a complemented disjoint sequence in $Y^r = c_0(\ell_{p_n}^{rM_n})$ for some $r \in \mathbb{N}$.

In this case our spaces X_n are modelled on the sets $J_n = \{1, 2, \dots, M_n\}$. Thus $c_0(\ell_{p_n}^{M_n})$ is modelled on the set $\{(n, k) : n \in \mathbb{N}, 1 \leq k \leq M_n\}$.

Now suppose $(u_n)_{n \in \mathbb{N}}$ is a C -tempered C -complemented disjoint sequence in $c_0(\ell_{p_n}^{M_n})$ with the property that $\|u_n[i]\|_{\ell_{p_i}^{M_i}} = 1$ or 0 for each i , and let $A = \{i : \exists n, u_n[i] \neq 0\}$. We claim that:

Claim: there exists a constant $K = K(C)$, an integer r depending only on C , a subset B of A with $|B| \leq r$ and $P_i \leq rM_i$ for $i \in B$ so that $(u_n)_{n \in \mathbb{N}}$ is K -equivalent to the canonical basis of $(\sum_{i \in B} \oplus \ell_{p_i}^{P_i})_{c_0}$.

To this end let $G(i) = \sup \langle u_n[i], u_n^*[i] \rangle$ as usual. We have $\sum_{i=1}^{\infty} G(i) \leq C$. We first consider the case when $G(i) \leq \frac{1}{20}$ for every i . Then it is possible to find a finite increasing sequence of integers $(k_j)_{j=0}^N$ where $N = [10C]$ depends on C such that

$$\sum_{i < k_j} G(i) \leq j/10$$

and

$$\sum_{i \leq k_j} G(i) \geq j/10.$$

Notice that

$$\sum_{i > k_N} G(i) \leq 1/10.$$

It follows that for each n there exist at least three values of $1 \leq j \leq N$ so that

$$\sum_{i=k_{j-1}+1}^{k_j} \langle u_n[i], u_n^*[i] \rangle > \frac{1}{4N}.$$

We can then assign to each n a value of j which is neither the largest or smallest with this property. In this way we partition $\mathcal{N}$ into sets $(\mathcal{N}_j)_{2 \leq j \leq N-1}$.

Consider $(u_n)_{n \in \mathcal{N}_j}$. We note that this is equivalent (with constants depending only on C) to each of $(v_n)_{n \in \mathcal{N}_j}$ and $(w_n)_{n \in \mathcal{N}_j}$ where $v_n[i] = u_n[i]$ if $i \leq k_{j-1}$ and 0 otherwise while $w_n[i] = u_n[i]$ if $i > k_j$ and 0 otherwise.

Now for any finitely non-zero sequence $(a_n)_{n \in \mathcal{N}_j}$ we have

$$\left\| \sum_{n \in \mathcal{N}_j} a_n v_n \right\|_Y \leq \left(\sum_{n \in \mathcal{N}_j} |a_n|^{q_{j-1}} \right)^{1/q_{j-1}}$$

where $q_j = p_{k_j}$. On the other hand

$$\left\| \sum_{n \in \mathcal{N}_j} a_n w_n \right\|_Y \geq \max_{i > k_j} \left(\sum_{u_n[i] \neq 0} |a_n|^{q_j} \right)^{1/q_j}.$$

Thus

$$\left\| \sum_{n \in \mathcal{N}_j} a_n w_n \right\|_Y^{q_j} \geq \frac{1}{C} \sum_{i > k_j} \sum_{u_n[i] \neq 0} G(i) |a_n|^{q_j}.$$

Now for fixed n

$$\sum_{i > k_j, u_n[i] \neq 0} G(i) \geq \sum_{i > k_j} \langle u_n[i], u_n^*[i] \rangle \geq \frac{1}{4N}.$$

It follows that $(u_n)_{n \in \mathcal{N}_j}$ satisfies an upper q_{j-1} -estimate and a lower q_j -estimate with constants depending only on C . We next estimate $|\mathcal{N}_j|$. In fact

$$\begin{aligned} \frac{1}{4N} |\mathcal{N}_j| &\leq \sum_{n \in \mathcal{N}_j} \sum_{i=k_{j-1}+1}^{k_j} \langle u_n[i], u_n^*[i] \rangle \\ &\leq \sum_{i=k_{j-1}+1}^{k_j} G(i) M_i \\ &\leq C \max_{k_{j-1} < i \leq k_j} M_i \end{aligned}$$

Hence if we select $k_{j-1} < i \leq k_j$ appropriately we have $(u_n)_{n \in \mathcal{N}_j}$ equivalent to a subset of the standard basis of $\ell_{p_i}^{rM_i}$ where r and the constant of equivalence depend only on C .

We must now treat the case when $G(i) > \frac{1}{20}$ for some i . In this case we split $\mathcal{N}$ into two groups $\mathcal{N}'$ and $\mathcal{N}''$ where $\mathcal{N}' = \{n : \exists i, \langle u_n[i], u_n^*[i] \rangle > \frac{1}{20}\}$ and $\mathcal{N}''$ is the remainder. Then $\mathcal{N}''$ can be treated as before. For $\mathcal{N}'$ we note that $(u_n)_{n \in \mathcal{N}'}$ is equivalent to a sequence (u'_n) where $u'_n[i] = u_n[i]$ for precisely one index $i = i_n$ such that $\langle u_n[i_n], u_n^*[i_n] \rangle > 1/20$ and is zero elsewhere. The

appropriate representation of $(u_n)_{n \in \mathcal{N}'}$ follows once we observe that the set $\{i_n : n \in \mathcal{N}'\}$ is bounded in cardinality with a bound depending only on C . But this is clear since $G(i_n) > 1/20$ but $\sum G(i) \leq C$.

Thus the claim is established.

Returning to our original hypotheses we see that if (u_n) is any unconditional basis of $c_0(\ell_{p_n}^{N_n})$ then (u_n) is equivalent to the canonical basis of $c_0(\ell_{p_n}^{M_n})$ where $M_n \leq rN_n$ for all n and some fixed r . By the same token the canonical basis of $c_0(\ell_{p_n}^{N_n})$ is equivalent a subset of $(u_n)^s$ for some $s \in \mathbb{N}$.

Now the additional hypotheses on N_n ensure that the original basis is equivalent to its square. Hence the s -fold product $(u_n)^s$ is equivalent to a subset of the canonical basis and so it follows from the Cantor-Bernstein principle (apparently first noticed by Mityagin, [14],[16] and [17]), that $(u_n)^s$ and the original basis are equivalent.

Thus the canonical bases of $c_0(\ell_{p_n}^{sM_n})$ and $c_0(\ell_{p_n}^{sN_n})$ are equivalent. Let $\mathcal{M} = \{(i, j) : 1 \leq j \leq M_i\}$ and $\mathcal{N} = \{(i, j) : 1 \leq j \leq N_i\}$. Let us suppose the former basis is indexed by $\mathcal{M}^s = \{(i, j) : 1 \leq j \leq sM_i\}$ and the latter by $\mathcal{N}^s = \{(i, j) : 1 \leq j \leq sN_i\}$. Let $\varphi : \mathcal{M}^s \rightarrow \mathcal{N}^s$ be a bijection implementing the claimed equivalence of bases. By elementary considerations concerning c_0 -sums it is clear that for each fixed i the set $\{\varphi(i, j) : 1 \leq j \leq M_i\}$ can have at most t distinct first co-ordinates where t depends only on the constant of equivalence; similarly for each fixed i the set of possible first co-ordinates of $\varphi^{-1}(i, j)$ can be bounded by the same t . For each $(a, b) \in \mathbb{N}^2$, let E_{ab} be the set of (i, j) so that $i = a$ and the first co-ordinate of $\varphi(i, j)$ is b . Then let F_{ab} be a subset of E_{ab} of size $\lfloor |E_{ab}|/s \rfloor$. Restricting φ to $\cup_{(a,b)} F_{ab}$ produces an equivalence between the bases of $c_0(\ell_{p_n}^{M_n - \alpha_n})$ and $c_0(\ell_{p_n}^{N_n - \beta_n})$ where $0 \leq \alpha_n, \beta_n \leq (s-1)t$ for each n . This clearly implies the equivalence of (u_n) and the original basis. $\square$

4. UNIQUENESS OF UNCONDITIONAL BASES IN c_0 -PRODUCTS OF RIGHT-DOMINANT SPACES

We first introduce some standard notation. Let A, B be subsets of $\mathbb{N}$. We write $A < B$ to indicate that $\max\{a : a \in A\} < \min\{b : b \in B\}$.

Let X be a sequence space modelled on $\mathbb{N}$. We say that X is *right-dominant* if there is a constant $\kappa = \kappa(X)$ so that whenever $u_1, \dots, u_n$ and $v_1, \dots, v_n$ are any two disjointly supported sequences satisfying $\text{supp } u_k < \text{supp } v_k$ and $\|u_k\|_X = \|v_k\|_X$ for $1 \leq k \leq n$ then $\|\sum_{k=1}^n u_k\|_X \leq \kappa \|\sum_{k=1}^n v_k\|_X$. We say that

X is *left-dominant* if there is a constant $\rho = \rho(X)$ so that whenever $u_1, \dots, u_n$ and $v_1, \dots, v_n$ are any two disjointly supported sequences satisfying $\text{supp } u_k < \text{supp } v_k$ and $\|u_k\|_X = \|v_k\|_X$ for $1 \leq k \leq n$ then $\|\sum_{k=1}^n v_k\|_X \leq \rho \|\sum_{k=1}^n u_k\|_X$.

Left and right-dominant spaces were studied in [4]. It is established (Lemma 5.2 of [4]) that in these spaces there is exactly one $r = r(X)$ (the *index* of X) so that ℓ_r is disjointly finitely representable in X . If X is right-dominant then X satisfies an upper r -estimate and a lower s -estimate for any $s > r$; the corresponding dual statements hold for left-dominant spaces. Clearly if a space X is both left and right-dominant then $X = \ell_r$.

Theorem 4.1. *Let X be a right-dominant sequence space with $r(X) = 1$. Then every complemented unconditional basic sequence in $c_0(X)$ is equivalent to a subsequence of the canonical basis.*

Remark. In particular this applies when X is a Nakano space $\ell_{(p_n)}$ where $p_n \downarrow 1$ or when X is Tsirelson space T (see [4]).

Proof. In this case we note that in the notation of Section 3, $J_i = \mathbb{N}$ for all $i \in \mathbb{N}$. We first note that by Corollary 2.5, $c_0(X)$ is not sufficiently Euclidean. Hence by [4] Theorem 3.5, every complemented unconditional basic sequence is equivalent to a complemented positive disjoint sequence in $c_0(X)^N$ for some N and hence also to a complemented disjoint sequence in $c_0(X)$.

Now by Theorem 3.2 it will suffice to show that if $(u_n)_{n \in \mathcal{N}}$ is a C -tempered C -complemented unconditional basic sequence then $(u_n)_{n \in \mathcal{N}}$ is K -equivalent to a subsequence of the canonical basis of $c_0(X)$ where K depends only on C . In fact we will show that it is K -equivalent to a subsequence of the canonical basis of $\ell_\infty^N(X)$ where N depends only on C .

We may suppose that, as before, $\|u_n[i]\|_X = 1$ or $u_n[i] = 0$. Let $f_n = u_n u_n^*$ and $F_n(i) = \langle u_n[i], u_n^*[i] \rangle$. Let $G(i) = \max_n F_n(i)$ so that $\|G\|_1 \leq C$.

Pick any integer $N > 2C$. For each $n \in \mathcal{N}$ we pick natural numbers $r_n \leq s_n$ such that

$$\sum_{i=1}^{\infty} \sum_{j=1}^{r_n-1} f_n(i, j) < \frac{1}{4}$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{r_n} f_n(i, j) \geq \frac{1}{4}$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{s_n-1} f_n(i, j) < \frac{3}{4}$$

$$\sum_{i=1}^{\infty} \sum_{j=1}^{s_n} f_n(i, j) \geq \frac{3}{4}.$$

We will argue by Hall's Marriage Lemma (see Bollobas [1]) that it is possible to find an map $\varphi : \mathcal{N} \rightarrow \mathbb{N}$ such that $\varphi(n) \in [r_n, s_n]$, with $|\varphi^{-1}(k)| \leq N$ for all $k \in \mathbb{N}$. Indeed if not, the Marriage Lemma implies there is a minimal finite subset $\mathcal{M}$ of $\mathcal{N}$ such that $N|\cup_{n \in \mathcal{M}} [r_n, s_n]| < |\mathcal{M}|$. It follows easily from the minimality that $\cup_{n \in \mathcal{M}} [r_n, s_n]$ is an interval $[a, b]$. From the disjointness of the (f_n) we have that

$$\sum_{j=a}^b \sup_{n \in \mathcal{M}} f_n(i, j) \leq (b - a + 1)G(i)$$

so that

$$\sum_{i=1}^{\infty} \sum_{j=a}^b f_n(i, j) \leq (b - a + 1)C$$

However

$$\sum_{i=1}^{\infty} \sum_{j=a}^b f_n(i, j) \geq \frac{1}{2}|\mathcal{M}|$$

so that $|\mathcal{M}| \leq 2(b - a + 1)C < N(b - a + 1)$ which is a contradiction.

We can now split $\mathcal{N}$ into at most N disjoint subsets $(\mathcal{N}_k)_{k \in \mathcal{M}}$ so that φ is injective on each $\mathcal{N}_k$.

For each $n \in \mathcal{N}$ let $A_n = \{(i, j) : j \leq \varphi(n)\}$ and $B_n = \{(i, j) : j \geq \varphi(n)\}$. Then, by Lemma 3.1, we have that $(u_n)_{n \in \mathcal{N}}$ is equivalent to both $(u_n \chi_{A_n})_{n \in \mathcal{N}}$ and $(u_n \chi_{B_n})_{n \in \mathcal{N}}$ with constants of equivalence depending only on C .

Now suppose $k \in \mathcal{M}$, and let $(a_n)_{n \in \mathcal{N}_k}$ be a finitely non-zero sequence. Then, by the right-dominance property, for each i

$$\left\| \sum_{n \in \mathcal{N}_k} a_n u_n \chi_{A_n} [i] \right\|_X \leq \kappa \left\| \sum_{n \in \mathcal{N}_k} a_n \|u_n \chi_{A_n} [i]\|_X e_{\varphi(n)} \right\|_X$$

Hence

$$(4.1) \quad \left\| \sum_{n \in \mathcal{N}_k} a_n u_n \chi_{A_n} [i] \right\|_Y \leq \kappa \left\| \sum_{n \in \mathcal{N}} a_n e_{\varphi(n)} \right\|_X.$$

In the opposite direction, again by the right-dominance property, we have

$$(4.2) \quad \left\| \sum_{n \in \mathcal{N}_k} a_n \|u_n \chi_{B_n}[i]\|_X e_{\varphi(n)} \right\|_X \leq \kappa \left\| \sum_{n \in \mathcal{N}_k} a_n u_n \chi_{B_n}[i] \right\|_X.$$

Combining (4.1) and (4.2),

$$(4.3) \quad \left\| \sum_{n \in \mathcal{N}_k} a_n \sum_{i=1}^{\infty} G(i) \|u_n \chi_{B_n}[i]\|_X e_{\varphi(n)} \right\|_X \leq C \kappa \left\| \sum_{n \in \mathcal{N}_k} a_n u_n \right\|_Y.$$

For each n let $I_n = \{i : 8CG(i) \geq \|u_n^*[i]\|_{X^*}\}$. Then

$$\sum_{i \notin I_n} G(i) \leq \frac{1}{8C} \sum_{i=1}^{\infty} \|u_n^*[i]\|_{X^*} \leq \frac{1}{8}.$$

Hence

$$\sum_{i \notin I_n} \langle u_n[i] \chi_{B_n}, u_n^*[i] \rangle \leq \frac{1}{8}.$$

However by choice of $\varphi(n)$ we have

$$\sum_{i=1}^{\infty} \langle u_n[i] \chi_{B_n}, u_n^*[i] \rangle \geq \frac{1}{4}.$$

Thus we have

$$\begin{aligned} \frac{1}{8} &\leq \sum_{i \in I_n} \|u_n[i] \chi_{B_n}\|_X \|u_n^*[i]\|_{X^*} \\ &\leq 8C \sum_{i \in I_n} G(i) \|u_n[i] \chi_{B_n}\|_X \\ &\leq 8C \sum_{i=1}^{\infty} G(i) \|u_n[i] \chi_{B_n}\|_X. \end{aligned}$$

The estimate above combined with (4.3) yields the inequality

$$\left\| \sum_{n \in \mathcal{N}_k} a_n e_{\varphi(n)} \right\|_X \leq 64C^2 \kappa \left\| \sum_{n \in \mathcal{N}_k} a_n u_n \right\|_Y.$$

Thus each $(u_n)_{n \in \mathcal{N}_k}$ is equivalent to $(e_{\varphi(n)})_{n \in \mathcal{N}_k}$ in X with constant of equivalence depending only on C . Since $|\mathcal{M}| \leq N$ where N depends only on C , the result is proved. $\square$

Let us say that an unconditional basis $(u_n)_{n \in \mathcal{N}}$ is *molecular* if there exists a constant C and a natural number N so that if $\mathcal{N}$ is partitioned into N disjoint sets $(\mathcal{N}_k)_{k=1}^N$ then there exists a proper subset $\mathcal{M}$ of $\{1, 2, \dots, N\}$ such that $(u_n)_{n \in \mathcal{N}}$ is C -equivalent to a subset of $\cup_{k \in \mathcal{M}} (u_n)_{n \in \mathcal{N}_k}$. Otherwise we will say that $(u_n)_{n \in \mathcal{N}}$ is *non-molecular*. It follows from the quantitative form of the Cantor-Bernstein principle [14,16,17] that $(u_n)_{n \in \mathcal{N}}$ is molecular if and only if there is a constant C so that if $\mathcal{N}$ is partitioned into N disjoint sets $(\mathcal{N}_k)_{k=1}^N$ then there is a proper subset $\mathcal{M}$ of $\{1, 2, \dots, N\}$ so that $(u_n)_{n \in \mathcal{N}_k, k \in \mathcal{M}}$ is C -equivalent to $(u_n)_{n \in \mathcal{N}}$. Let us note that any subsymmetric basis is molecular with $N = 2$ as is the usual basis of $(\sum_{n=1}^{\infty} \ell_q^n)_{\ell_p}$ when $1 \leq p, q < \infty$. The canonical basis of $\ell_p \oplus \ell_q$ for $p \neq q$ is molecular with $N = 3$.

Lemma 4.2. *Suppose $(u_n)_{n=1}^{\infty}$ is a non-molecular unconditional basis. Then for any $\epsilon > 0$, $N \in \mathbb{N}$ and constant C there exists $M > N$ and subsets $(A_k)_{k=1}^M$ of $\mathbb{N}$ so that:*

- (1) *For any subset $\mathcal{M}$ of $\{1, 2, \dots, M\}$ with $|\mathcal{M}| < N$ then $(u_n)_{n=1}^{\infty}$ is not C -equivalent to any subset of $\{u_n : n \in \cup_{k \in \mathcal{M}} A_k\}$, and*
- (2) $\frac{1}{M} \sum_{k=1}^M \chi_{A_k} \geq (1 - \epsilon) \chi_{\mathbb{N}}$.

Proof. It suffices to consider the case when $\epsilon = r/s$ is rational. We then may pick an integer m so large that $mr > N$ and so that if $L = \binom{ms}{mr}$ then we can partition $\mathbb{N}$ into L sets so that (u_n) is not C -equivalent to a subset of $(u_n)_{n \in \mathcal{N}}$ where $\mathcal{N}$ is the union of any $L - 1$ sets.

Let Ω be the collection of all $m(s-r)$ subsets of $\{1, 2, \dots, ms\}$. We can partition $\mathbb{N} = \cup_{\omega \in \Omega} B_{\omega}$ so that $(u_n)_{n \in \mathbb{N}}$ is not C -equivalent to a subset of $(u_n)_{n \in \mathcal{N}}$ where $\mathcal{N} = \cup_{\omega \in D} B_{\omega}$ for some proper subset D of Ω .

Now let $A_k = \cup_{\omega: k \in \omega} B_{\omega}$ for $1 \leq k \leq M = ms$. It is clear that

$$\sum_{k=1}^{ms} \chi_{A_k} = m(s-r) \chi_{\mathbb{N}}$$

so that (2) holds. Suppose $(u_n)_{n \in \mathbb{N}}$ is C -equivalent to a subset of $(u_n)_{n \in \mathcal{N}}$ where $\mathcal{N} = \cup_{k \in \mathcal{M}} A_k$. Then we have $\cup_{k \in \mathcal{M}} A_k = \Omega$ whence $|\mathcal{M}| > mr$. $\square$

Theorem 4.3. *Let X be a space with nontrivial cotype and an unconditional basis (u_n) . If $c_0(X)$ has a unique unconditional basis then (u_n) is molecular.*

Proof. We will assume, on the contrary that the basis (u_n) is not molecular. Let us regard X as a sequence space so that the given unconditional basis is

identified with $(e_n)_{n \in \mathbb{N}}$. We start by using Lemma 4.2 repeatedly to generate for each $r \in \mathbb{N}$, subsets $(A_{rk})_{k=1}^{M_r}$ of $\mathbb{N}$ so that:

- (1) For any subset $\mathcal{M}$ of $\{1, 2, \dots, M_r\}$ with $|\mathcal{M}| < r$ the basis $(e_n)_{n=1}^\infty$ is not r -equivalent to any subset of $\{e_n : n \in \cup_{k \in \mathcal{M}} A_{rk}\}$, and
- (2) $M_r^{-1} \sum_{k=1}^{M_r} \chi_{A_{rk}} \geq (1 - 2^{-(r+1)}) \chi_{\mathbb{N}}$.

Now for each $s \in \mathbb{N}$ let $P_s = \prod_{r=1}^s M_r$ and let $(B_{sk})_{k=1}^{P_s}$ be a listing of all sets of the form $\cap_{r=1}^s A_{rk_r}$. We observe that

$$P_s^{-1} \sum_{k=1}^{P_s} \chi_{B_{sk}} \geq \frac{1}{2} \chi_{\mathbb{N}}.$$

Consider the index set $I = \{(s, k) : 1 \leq k \leq P_s, s \in \mathbb{N}\}$. We will consider the space $c_0(X)$ as a sequence space modelled on $I \times \mathbb{N}$.

Consider now the block basic sequence

$$u_{sn} = \sum_{k=1}^{P_s} e_{skn}.$$

If we define the biorthogonal functionals

$$u_{sn}^* = \frac{1}{P_s} \sum_{k=1}^{P_s} e_{skn}$$

then it is clear that $(u_{sn})_{s,n}$ is a complemented disjoint sequence equivalent to the canonical basis of $c_0(X)$.

Now let $D = \{(s, k, n) : n \in B_{sk}\}$. Then $\|u_n u_n^* \chi_D\|_1 \geq \frac{1}{2}$. It follows from Lemma 3.1 that $(u_{sn} \chi_D)_{s,n}$ is also a complemented disjoint sequence equivalent to the canonical basis of $c_0(X)$.

The basis vectors (e_{skn}) for $(s, k, n) \in D$ span a complemented subspace Y of $c_0(X)$ which by the above remark contains a complemented copy of $c_0(X)$. By the Pełczyński decomposition argument, Y is isomorphic to $c_0(X)$. If we assume that $c_0(X)$ has a unique unconditional basis then it will follow that the whole basis $(e_{skn})_{(s,k,n) \in I, n \in \mathbb{N}}$ is C -equivalent, for some C , to its subset $(e_{skn})_{(s,k,n) \in D}$.

Thus we can partition D into subsets $(D_t)_{t=1}^\infty$ so that each subset (e_{skn}) for $(s, k, n) \in D_t$ is C -equivalent to the canonical basis (e_n) of X while any subset obtained by picking one element from each D_t is C -equivalent to the standard c_0 -basis. From this and the fact that X has a lower-estimate it is clear that

for fixed (s, k) at most finitely many D_t can intersect the set of all (s, k, n) for $n \in \mathbb{N}$. Note also that the set of (s, k) such that $(s, k, n) \in D_t$ for some n must also be uniformly bounded by some constant K again by the lower estimate on X .

In particular for any s_0 there exists t so that if $(s, k, n) \in D_t$ then $s > s_0$. Hence, the canonical basis of X is C -equivalent to a subset of $\cup_{(s,k) \in \mathcal{M}} B_{sk}$ where $(s, k) \in \mathcal{M}$ implies $s > s_0$ and $|\mathcal{M}| \leq K$. Now each B_{sk} is contained in some $A_{s_0, k}$ and so we must have $K \geq s_0$. By choosing s_0 large enough we get a contradiction. $\square$

We now state without proof a general theorem which can be proved by exactly the same argument.

Theorem 4.4. *Suppose $1 \leq p < \infty$ and suppose X is a Banach space with a non-molecular unconditional basis $(u_n)_{n=1}^\infty$ with the property that it does not contain subsets uniformly equivalent to the unit vector bases of ℓ_p^m for $m = 1, 2, \dots$. Let $(u_{mn})_{m,n=1}^\infty$ be the induced basis of $\ell_p(X)$. Then there is a subset $\mathcal{A} \subset \mathbb{N} \times \mathbb{N}$ so that $(u_{mn})_{(m,n) \in \mathcal{A}}$ is non-equivalent to the full basis (u_{mn}) and spans a subspace isomorphic to $\ell_p(X)$.*

We conclude with a theorem which gives us a large number of examples of right-dominant spaces with non-molecular unconditional bases.

Theorem 4.5. *Suppose X is a right-dominant sequence space with $r(X) = r$. Suppose the canonical basis is molecular. Then $X = \ell_r$.*

Proof. It is enough to show that X is left-dominant. Let us assume the contrary. Then we claim:

Claim: Given any $a \in \mathbb{N}$ and $C > 0$ there exists $b > a$ so that $(e_k)_{a < k \leq b}$ is not C -equivalent to any subset of $(e_k)_{k \leq a} \cup (e_k)_{b < k}$.

To prove the claim let $C_1 > C^2\kappa + a$. Since X is not left-dominant there exist disjoint sequences $(u_n)_{n=1}^N$ and $(v_n)_{n=1}^N$ with finite supports so that $a < \text{supp } u_n < \text{supp } v_n$ for each n , $\|u_n\|_X = \|v_n\|_X$ and

$$\left\| \sum_{n=1}^N v_n \right\|_X > C_1 \left\| \sum_{n=1}^N u_n \right\|_X.$$

Pick b so large that $\text{supp } v_n \leq b$ for all n . Suppose $(e_k)_{a < k \leq b}$ is C -equivalent to some subset of $(e_k)_{k \leq a} \cup (e_k)_{b < k}$. Then there exist $(w_n)_{n=1}^N$ each with finite

disjoint supports not intersecting $(a, b]$ so that $\|w_n\|_X = \|u_n\|_X$ for $1 \leq n \leq N$ and

$$\left\| \sum_{n=1}^N w_n \right\|_X \leq C^2 \left\| \sum_{n=1}^N u_n \right\|_X.$$

Let $\mathcal{M} = \{n : \text{supp } w_n \cap [1, a] \neq \emptyset\}$. Then $|\mathcal{M}| \leq a$. Thus

$$\left\| \sum_{n \in \mathcal{M}} v_n \right\|_X \leq a \left\| \sum_{n=1}^N u_n \right\|_X.$$

On the other hand

$$\left\| \sum_{n \notin \mathcal{M}} v_n \right\|_X \leq \kappa \left\| \sum_{n \notin \mathcal{M}} w_n \right\|_X \leq C^2 \kappa \left\| \sum_{n=1}^N u_n \right\|_X.$$

It follows that $C_1 < C^2 \kappa + a$ contrary to assumption. This establishes the claim.

To prove the theorem we use the claim to find an increasing sequence $(a_n)_{n=1}^\infty$ so that $(e_k)_{a_n < k \leq a_{n+1}}$ is not n -equivalent to any subset of $(e_k)_{k \leq a_n} \cup (e_k)_{k > a_{n+1}}$. Then fix any $s \in \mathbb{N}$ and consider the sets $A_j = \cup \{(a_n, a_{n+1}] : n \equiv j \pmod s\}$ for $0 \leq j \leq s-1$. Now the sets $(e_n)_{n \in A_j}$ partition the basis into s sets in such a way that no $(s-1)$ sets contain a subset equivalent to the original basis. This contradicts our assumption that the basis is molecular. $\square$

Examples. We can now give many examples of spaces X with a unique unconditional basis but such that the c_0 -product $c_0(X)$ fails to have unique unconditional basis. This will answer negatively a question raised in [3].

In fact if X is right-dominant and $c_0(X)$ has unique unconditional basis then X must be one of the three spaces c_0, ℓ_1 or ℓ_2 . This follows by observing that if it is not in this list then $r(X) < \infty$ and hence X has cotype. Then the preceding Theorems 4.3 and 4.5 show that $X = \ell_r$ for some finite r . The uniqueness then forces either $r = 1$ or $r = 2$.

On the other hand there are many known examples of right-dominant spaces with unique unconditional bases. In [3] 2-convexified Tsirelson space is shown to have unique unconditional bases. In [4] Tsirelson space itself and certain Nakano spaces $\ell_{(p_n)}$ are shown to have unique unconditional bases. These latter examples satisfy $r(X) = 1$ so that we can apply Theorem 4.1. The second non-equivalent basis constructed in Theorem 4.3 is indeed equivalent to a subset of the original basis. $\square$

REFERENCES

1. B. Bollobas, *Combinatorics*, Cambridge University Press, 1986.
2. J. Bourgain, *On the Dunford-Pettis property*, Proc. Amer. Math. Soc. **81** (1981), 265-272.
3. J. Bourgain, P.G. Casazza, J. Lindenstrauss and L. Tzafriri, *Banach spaces with a unique unconditional basis, up to a permutation*, Memoirs Amer. Math. Soc. No. 322, 1985.
4. P.G. Casazza and N.J. Kalton, *Uniqueness of unconditional bases in Banach spaces*, Israel J. Math **103** (1998), 141-176.
5. P.G. Casazza and T.J. Schura, *Tsirelson's space*, Springer Lecture Notes 1363, 1989.
6. I.S. Edelstein and P. Wojtaszczyk, *On projections and unconditional bases in direct sums of Banach spaces*, Studia Math. **56** (1976), 263-276.
7. T. Figiel, J. Lindenstrauss and V.D. Milman, *The dimension of almost spherical sections of convex bodies*, Acta Math. **139** (1977), 53-94.
8. W.T. Gowers, *A solution to Banach's hyperplane problem*, Bull. London Math. Soc. **26** (1994), 523-530.
9. G. Köthe and O. Toeplitz, *Lineare Räume mit unendlich vielen Koordinaten und Ringen unendlicher Matrizen*, J. Reine Angew. Math. **171** (1934), 193-226.
10. J. Lindenstrauss and A. Pełczynski, *Absolutely summing operators in $\mathcal{L}_p$ -spaces and their applications*, Studia Math. **29** (1968), 315-349.
11. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I, Sequence spaces*, Springer Verlag, Berlin, Heidelberg, New York, 1977.
12. J. Lindenstrauss and M. Zippin, *Banach spaces with a unique unconditional basis*, J. Functional Analysis **3** (1969), 115-125.
13. V.D. Milman and G. Schechtman, *Asymptotic theory of finite-dimensional spaces*, Springer Lecture Notes 1200, 1986.
14. B.S. Mityagin, *Equivalence of bases in Hilbert scales (in Russian)*, Studia Math. **37** (1970), 111-137.
15. G. Pisier, *The volume of convex bodies and geometry of Banach spaces*, Cambridge Tracts 94, Cambridge University Press, 1989.
16. P. Wojtaszczyk, *Uniqueness of unconditional bases in quasi-Banach spaces with applications to Hardy spaces, II*, Israel J. Math **97** (1997), 253-280.
17. M. Wojtowicz, *On Cantor-Bernstein type theorems in Riesz spaces*, Indag. Math. **91** (1988), 93-100.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO. 65211, U.S.A.

E-mail address: `pete@math.missouri.edu`, `nigel@math.missouri.edu`